

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 1
		Version 4.0
		15. August 2023

ESA Climate Change Initiative “Plus” (CCI+)

Product User Guide (PUG) Version 4.0

-

for the RemoTeC XCO₂ GOSAT-2 SRON Full-Physics Product (CO₂_GO₂_SRFP) Version 2.0.2

for the Essential Climate Variable (ECV)

Greenhouse Gases (GHG)

Written by:

GHG-CCI+ project team

Lead author:

Andrew Gerald Barr

(SRON Netherlands Institute for Space Research, Leiden, The Netherlands)

Co-authors:

Tobias Borsdorff, Jochen Landgraf

(SRON Netherlands Institute for Space Research, Leiden, The Netherlands)

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 2
		Version 4.0
		15. August 2023

Change log:

Version Nr.	Date	Status	Reason for change
Version 1	27. Oct. 2020	Draft	New document
Version 1.1	04. Jan. 2021	As submitted	- Update format - Update purpose of document
Version 1.1	04. Feb. 2021	As submitted	- Update after ESA reviews - Remote typos
Version 2.0	04. Nov. 2021	As submitted	- L2 data reprocessing: update filter criteria, selection of TCCON station, and bias correction
Version 3.0	27. Jan. 2022	As submitted	- Updated doc to version 3.0
Version 4.0	15. Aug. 2023	As submitted	- Update doc to version 4.0 - Quality filtering via random forest model prediction

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 3
		Version 4.0
		15. August 2023

Table of Contents

1.	Purpose of document	4
2.	Greenhouse gases Observing SATellite-2 (GOSAT-2)	5
3.	RemoTeC retrieval algorithm.....	6
4.	XCO ₂ RemoTeC FP data product (Feb. 2019 – Dec 2021).....	7
	4.1 Global maps.....	7
	4.2 Validation with TCCON	10
	4.3 Bias correction	16
5.	Description of data format	16
	5.1 Product Content and Format.....	16
	5.2 Quality Flags and Metadata	20
	5.3 Recommended data usage	24
	5.4 Tools for Reading the Data	25
6.	References	26

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 4
		Version 4.0
		15. August 2023

1. Purpose of document

This document describes the Product User Guide (PUG) of the RemoTeC XCO₂ GOSAT-2 SRON Full-Physics Product (CO₂_GO₂_SRFP), which is a deliverable for the ESA GHG-CCI+ project led by University of Bremen, Germany.

Within the project, satellite-derived atmospheric Carbon Dioxide (CO₂) and Methane (CH₄) Essential Climate Variable (ECV) data products are generated and delivered to ESA for inclusion into the ESA-GHG-CCI+ database from which users can access these data products and the corresponding documentations.

The satellite-derived data products are:

- Column-averaged dry-air mixing ratios (mole fractions) of CO₂ and CH₄, denoted XCO₂ (in parts per million, ppm) and XCH₄ (in parts per billion, ppb), respectively.

This document will be focused on the XCO₂ Level-2 product retrieved using the GOSAT-2 Full-Physics algorithm developed by SRON Netherlands Institute for Space Research, The Netherlands.

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 5
		Version 4.0
		15. August 2023

2. Greenhouse gases Observing SATellite-2 (GOSAT-2)

The Japanese Greenhouse gases Observing SATellite-2 (GOSAT2) was launched on 29th October 2018 and started operational observations from February 2019. GOSAT2 provides dedicated global measurements of total column CO₂ and CH₄ from its SWIR bands. It is equipped with two instruments, the Thermal And Near Infrared Sensor for carbon Observations - Fourier Transform Spectrometer-2 (TANSO-FTS2) as well as a dedicated Cloud and Aerosol Imager-2 (TANSO-CAI-2).

The TANSO-FTS2 instrument (Nakajima et al., 2017) has five spectral bands with a high spectral resolution 0.2 cm⁻¹. Three operate in the SWIR at 0.75-0.77, 1.56-1.69 and at the extended 1.92-2.33 µm range, providing sensitivity to the near-surface absorbers. The fourth and fifth channels operating in the thermal infrared between 5.5-8.4 and 8.4-14.3 µm providing mid-tropospheric sensitivity.

The measurement strategy of TANSO-FTS2 is optimized for the characterization of continental-scale sources and sinks. TANSO-FTS2 utilizes a pointing mirror to perform off-nadir measurements at the same location on each 6-day repeat cycle. The pointing mirror allows TANSO-FTS2 to observe up to ±35° across track and ±40° along-track. These measurements nominally consist of 5 across track points spaced ~160km apart with a ground footprint diameter of approximately 9.7 km and a 4 second exposure duration. The satellite has an intelligent pointing monitor camera which makes it possible to adjust the line of sight of the FTS to steer away from cloud contaminated areas. Whilst the majority of data is limited to measurements over land where the surface reflectance is high, TANSO-FTS2 also observes in sun-glint mode over the ocean.

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 6
		Version 4.0
		15. August 2023

3. RemoTeC retrieval algorithm

The CO₂_GO₂_SRFP product is retrieved from GOSAT-2 TANSO-FTS spectra using the RemoTeC algorithm, that has been developed jointly by SRON and Karlsruhe Institute of Technology (KIT). The algorithm retrieves simultaneously XCH₄ and XCO₂. For the retrieval, we analyze four spectral regions, the 0.77 µm oxygen band, two CO₂ bands at 1.61 and 2.06 µm, as well as a CH₄ band at 1.64 µm. Within the retrieval procedure, the sub-columns of CO₂ and CH₄ in different altitude layers are being retrieved. To obtain the column averaged dry air mixing ratios XCO₂ and XCH₄ the sub-columns are summed up to get the total column, which is divided by the dry-air columns obtained from ECMWF model data in combination with a surface elevation data base.

The retrieved XCO₂ has been validated against ground based measurements from the Total Column Carbon Observing Network (TCCON). To further improve accuracy of XCO₂ product, a bias correction has been developed based on TCCON comparisons. We use the GGG2020 release of the TCCON data (Wunch et al., 2015, Laughner et al. 2021). More details on the technical aspects of the retrievals can be found in the ATBD GO₂-SRFP document (Barr et al. 2023a).

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 7
		Version 4.0
		15. August 2023

4. XCO₂ RemoTeC FP data product (Feb. 2019 – Dec 2021)

In this section, we show examples of the GOSAT-2 XCO₂ FP data product by showing global averaged maps (Sec. 4.1) and by giving a summary of the validation results relative to TCCON (Sec. 4.2).

4.1 Global maps

Figure 1 and Figure 2 show global average maps of the RemoTeC GOSAT-2 FP XCO₂ data product. Figure 1 shows the bias-corrected XCO₂ data and Figure 2 is the scaled random error (Barr et al. 2023b). The GOSAT-2 XCO₂ product provides good global spatial coverage. As can be seen, in some regions the coverage is limited by cloud cover (the observations correspond to cloud free scenes), sun illumination conditions, etc. The error on XCO₂ varies from 0.31 ppm to 7.4 ppm and is higher for high latitude regions and low albedo areas (Figure 2).

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO2_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 8
		Version 4.0
		15. August 2023




Figure 1: Global averaged XCO₂ (in ppm) between February 2019 and Dec 2021 for the CO₂_GO2_SRFP product on a 0.5x0.5° latitude/longitude grid (top) and 2x2° grid (bottom).

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO2_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 9
		Version 4.0
		15. August 2023




Figure 2: Same as Figure 1 but for the corresponding error on XCO₂ in ppm.

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 10
		Version 4.0
		15. August 2023

4.2 Validation with TCCON

This section summarizes the main validation results presented in the RemoTeC GOSAT-2 ESA GHG CCI+ End-to-End ECV Uncertainty Budget (E3UB) version 4.0 document (Barr et al. 2023b). We used ground based TCCON GGG2020 (Laughner et al., 2021) data obtained from <https://tccondata.org/> as reference data set. We co-located GOSAT-2 and TCCON measurements with a maximum time difference of 2.5h, a maximum distance of 300 km in both longitudinal and latitudinal directions. In cases of multiple TCCON measurements of the same site collocating with a GOSAT-2 sounding, we averaged the TCCON measurements.

The mean bias (global offset) amounts to -0.01 ppm. The standard deviation of the site biases (spatial accuracy or station-to-station variability) is 0.5 ppm for land and 1.31 ppm for ocean observations. TCCON observes these gases with a precision on mole fractions of ~0.15% and ~0.2% for CO₂ and CH₄, respectively (Toon et al., 2009). The single measurement precision of GOSAT-2 compared to TCCON amounts to 2.21 ppm. The validation results are summarised in Table 1.

Figure 3 shows the collocations of GOSAT-2 FP XCO₂ for land observations, and Figure 4 the same for observations over ocean (sun glint), with the TCCON sites. Statistics per site for each mode are shown in Figure 5 and Figure 6, respectively. Detailed bias and scatter (i.e., single sounding precision measured by the standard deviation of the difference to TCCON after removing systematic effects) are described in the E3UB (Barr et al. 2023b).

 GHG-CCI+ project	ESA Climate Change Initiative "Plus" (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO2_GO2_SRF) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 11
		Version 4.0
		15. August 2023

	Full Physics					
Variable	N	μ (ppm)	σ (ppm)	$\bar{\mu} \pm \sigma_{\bar{\mu}}$ (ppm)	$\bar{\sigma} \pm \sigma_{\bar{\sigma}}$ (ppm)	R
GOSAT2 Land	12557	-0.01	2.21	0.06 ± 0.50	2.28 ± 0.56	0.82
GOSAT-2 Ocean	118	-0.02	2.59	0.54 ± 1.31	2.39 ± 0.48	0.63

Table 1: Overview of the GOSAT-2 XCO₂ products vs TCCON co-located measurements. The mean bias μ and single measurement precision σ are calculated by taking the mean and standard deviation of the differences of all GOSAT-2 and TCCON pairs. The mean of the site means $\bar{\mu}$ and the spatial accuracy $\sigma_{\bar{\mu}}$ are calculated by taking the mean and standard deviation of the site means. The mean standard deviation $\bar{\sigma}$ and standard deviation of the standard deviations $\sigma_{\bar{\sigma}}$ are calculated by taking the mean and the standard deviation of the site standard deviations.

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO2_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 12
		Version 4.0
		15. August 2023




Figure 3: Validation of land soundings of XCO₂ with collocated TCCON measurements at all TCCON sites for the period Feb. 2019 - Dec 2021. Numbers in the figures: μ = bias, i.e., average of the difference; σ = single measurement precision, i.e., standard deviation of the difference; N = number of co-locations; R = Pearson correlation coefficient.




Figure 4: Validation of ocean soundings of XCO₂ with collocated TCCON measurements at all TCCON sites for the period Feb. 2019 - Dec 2021. Numbers in the figures: μ = bias, i.e., average of the difference; σ = single measurement precision, i.e., standard deviation of the difference; N = number of co-locations; R = Pearson correlation coefficient.




Figure 5: Validation statistics bias (top) and scatter (bottom) per TCCON site for land observations (bias corrected). The summarizing values represent the standard deviation of the site biases and the average scatter relative to TCCON.




Figure 6: Same as Figure 5 but for sun-glint observations.

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 16
		Version 4.0
		15. August 2023

4.3 Bias correction

From comparison with TCCON it was found, that the error in XCO₂ is highly correlated with the retrieved albedo α at window 2 (1600 nm). Based on this correlation, the following bias correction for land observations has been developed.

$$XCO_{2\text{corr}} = XCO_2 * (a + b * \alpha)$$

with $a = 0.99023$, $b = 0.05021$. The bias correction parameters are obtained by the fitting of GOSAT-2 and TCCON differences.

For sun-glint observations, it is found that XCO₂ error is correlated with the O₂ ratio RO₂. It defines the ratio between retrieved and prior O₂ column. In this case, a similar correction function is applied,

$$XCO_{2\text{corr}} = XCO_2 * (a + b * RO_2)$$

with $a = 1.46845$ and $b = -0.47389$.

5. Description of data format

5.1 Product Content and Format

The CO₂_GO₂_SRFP data products are stored per day in a single NetCDF file. Retrieval results are provided for the individual GOSAT-2 spatial footprints, i.e. no averaging has been applied. The product file contains the key standard products, i.e. the retrieved column averaged dry air mixing ratios XCO₂ and XCH₄ with bias correction, averaging kernels and quality flags, as well as secondary products specific for the RemoTeC algorithm. Tables 2-4 list the key variables in the data product.

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 17
		Version 4.0
		15. August 2023

Table 2: Common dimensions for the CO₂_GO₂_SRFP product.

Dimension	Type	Unlimited	Units	Description
sounding_dim	int	no		Number of sounding
polarization_dim	int	no		Number of polarization = 2
level_dim	int	no		Number of level = 13
layer_dim	int	no		Number of layer = 12
window_dim	int	no		Number of retrieval window = 4
char_l1bname	int	no		Number of character of L1B name = 44

Table 3: Common variables for the CO₂_GO₂_SRFP product.

Name	Type	Dim.	Units	Description
solar_zenith_angle	float	n	degrees	Angle between line of sight to the sun and local vertical
sensor_zenith_angle	float	n	degrees	Angle between the line of sight to the sensor and the local vertical
time	float	n	seconds	Seconds since 1970-01-01 00:00:00
longitude	float	n	degrees _east	Center longitude
latitude	float	n	degrees _north	Center latitude
pressure_levels	float	n, 13	hPa	Pressure levels
pressure_weight	float	n, 12		Layer dependent weights needed to apply the averaging kernels
xco2	float	n	1e-6	Retrieved column dry-air mole fraction of atmospheric carbon dioxide (XCO ₂) in ppm
xco2_uncertainty	float	n	1e-6	1-sigma uncertainty of the retrieved column-average dry-air mole fraction of atmospheric carbon dioxide

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 18
		Version 4.0
		15. August 2023

xco2_averaging_kernel	float	n, 12		Normalized column averaging kernel
co2_profile_apriori	float	n, 12	1e-6	A priori dry-air mole fraction profile of atmospheric carbon dioxide
xco2_quality_flag	int	n		Quality flag for XCO ₂ retrieval, 0 = good, 1 = bad

Table 4: Product specific (additional) variables for the CO₂_GO₂_SRFP product.

Name	Type	Dim.	Units	Description
flag_landtype	int	n		0 = land, 1 = ocean
flag_sunglint	int	n		0 = no sunglint, 1 = sunglint
gain	char	n		Number of gain coefficient calculated from solar calibration mode data. [1P 1S 2P 2S 3P 3S]
exposure_id	int	n		Exposure identification number of the sounding
l1b_name	char	n		Name of the Level 1B file of the sounding
signal_to_noise_window	float	n, 4, 2		Signal to noise ratio per retrieval window and for both polarization directions
dry_airmass_layer	float	n, 12	m-2	Dry airmass per layer
altitude	float	n	m	Vertical distance above the surface
air_temperature	float	n, 13	K	The bulk temperature of the air at each level
surface_elevation_stdev	float	n	m	Standard deviation of the surface elevation within the sounding
x_wind	float	n, 13	m s-1	Eastward wind velocity
y_wind	float	n, 13	m s-1	Northward wind velocity
chi2	float	n		Chi-squared value of the sounding
optical_thickness_of_atmosphere_layer_due_to_ambient_aerosol	float	n, 4		Scattering optical thickness per retrieval window

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 19
		Version 4.0
		15. August 2023

raw_xco2	float	n	1e-6	Retrieved column dry-air mole fraction of atmospheric carbon dioxide (XCO ₂) in ppm before bias correction
raw_xco2_err	float	n	1e-6	1-sigma statistical uncertainty of the retrieved column-average dry-air mole fraction of atmospheric carbon dioxide
h2o_column	float	n	m-2	Retrieved total water column
surface_albedo_758	float	n		The retrieved albedo at 758 nm
surface_albedo_1593	float	n		The retrieved albedo at 1593 nm
surface_albedo_1629	float	n		The retrieved albedo at 1629 nm
surface_albedo_2042	float	n		The retrieved albedo at 2042 nm
intensity_offset_o2a	float	n	W cm-2	The retrieved intensity offset in the O ₂ A band
aerosol_size	float	n		Retrieved size parameter of the aerosol distribution
aerosol_central_height	float	n	m	Peak height of the aerosol Gaussian height distribution
aerosol_total_column	float	n	m-2	Retrieved total aerosol column

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 20
		Version 4.0
		15. August 2023

5.2 Quality Flags and Metadata

To use data of GOSAT-2 FP XCO₂, users are encouraged to check the corresponding quality flag. In the NetCDF files, the quality flag, namely xco2_quality_flag, has been generated. It can have two values,

- 0: good quality data (quality has been checked)
- 1: data should not be used (e.g. bad fit to data, residual cloud contamination)

Filtering

In v2.0.2 of the XCO₂ full physics product, quality filtering is conducted through the use of a trained random forest classifier model which predicts the quality of the retrievals based on a selection of retrieval parameters, such as the cirrus signal, intensity offset, slope of the continuum etc.

The random forest model is trained on GOSAT-2 collocations with TCCON, where retrievals are classified via the bias of XCO₂. Retrievals in the training set are flagged as good quality if the bias is within a certain range and those outwith this range are flagged as bad quality. The model then learns the relationship between the quality of the retrieval and the selection of retrieval parameters on which it is trained, and uses these to predict the quality of all future data.

XCO₂ from TCCON is also used in the validation of the final product (sections 4.2 and 4.3). The random forest model is trained in a supervised way, where the model is given values for a selection of features as well as a classification label. Thus inclusion of the same data in both training and predicting can lead to artificial features, as the model has already seen the target label in training and therefore simply predicts this. In order to avoid this, we implement a temporal extrapolation by excluding one year from the training data, and using the resulting model to make the quality predictions for the excluded year. This results in one model per year of filtered data. Here we assume that the relationship between retrieval quality and features used in training is not temporally dependent. Figure 7 shows the classification metrics for the random forest model for 2021.




Figure 7: *Top:* The ROC curve (left) shows the true positive rate vs. false positive rate at different classification thresholds. Lowering the classification threshold classifies more items as positive, thus increasing both false positives and true positives. In the precision-recall (right) curve, a high area under the curve represents both high recall and high precision, where high precision relates to a low false positive rate, and high recall relates to a low false negative rate. *Bottom left:* List of features used in the model in order of importance from top to bottom. *Bottom right:* Confusion matrix comparing number of correct predictions.

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 22
		Version 4.0
		15. August 2023

A consequence of training the random forest model on GOSAT-2 collocations with TCCON is that retrievals with surface albedo ≥ 0.4 are absent from the training sample, due to the lack of TCCON stations in high albedo areas. To circumvent this we define a set of high albedo data to include in the training set using a list of threshold criteria (following v2.0.0 of this product). Furthermore the random forest method is limited only to land retrievals due to the low number of collocations over ocean. For ocean measurements we then also apply the threshold criteria for determining the quality flag. The criteria are as follows where any retrieval that does not satisfy all of these is assigned a 1 for the quality flag:

For land:

- Cost function (χ^2) < 12.0
- Number of iterations < 31
- Signal to noise ratio (SNR) > 50
- Std. deviation of surface elevation within GOSAT-2 ground pixel < 100 m
- Solar zenith angle (SZA) < 75°
- Aerosol optical thickness (AOT) window 1 < 1.0
- 3 < Aerosol size < 6
- 0 < Aerosol height < 10,000 m
- 0 < Blended Albedo < 1.4
- 0 < Cirrus signal < 2.0E-9
- 0.99 < CO₂ ratio < 1.018
- 0.96 < O₂ ratio < 1.04
- 0.95 < H₂O ratio < 1.08

For ocean (sun-glint):

- Cost function (χ^2) < 12.0
- Number of iterations < 31
- Signal to noise ratio (SNR) > 50

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 23
		Version 4.0
		15. August 2023

- Std. deviation of surface elevation within GOSAT-2 ground pixel < 100 m
- Solar zenith angle (SZA) < 75°
- 0 < Blended Albedo < 0.4
- 0 < Cirrus signal < 2.0E-9
- 0.99 < CO₂ ratio < 1.003
- 0.96 < O₂ ratio < 1.04
- 0.95 < H₂O ratio < 1.08

The yield and accuracy of the data of the final product is directly correlated to the strictness of the classification of the training dataset. For the data product described here, filtering using the random forest classifier results in an increase in 9 % for both data yield and precision in comparison to v2.0.0. The machine learning approach better captures the inter-dependencies of the retrieval parameters as it learns these during training, and thus less retrievals are filtered out when they should be kept in and vice versa.

Finally, we find an excellent correlation ($R=0.76$) and low bias (0.18 ppm) between v2.0.0 and v2.0.2 for land retrievals, illustrated in Figure 8, meaning that, while TCCON stations are limited in global coverage, training on TCCON data does not produce obvious biases when extrapolating the filtering to global scales.

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO2_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 24
		Version 4.0
		15. August 2023




Figure 8: Correlation plot of XCO₂ from v2.0.2 against v2.0.0. Data are sampled on a 1x1o grid across all latitude/longitudes for land measurements over the time period Feb 2019-Dec 2021.

5.3 Recommended data usage

It is strongly recommended to only use the bias-corrected data in, except if users explicitly correct for biases themselves (e.g. in an inverse modeling framework). The bias correction has been developed independently for land and sun-glint observations.

Important! Please only use data with xco2_quality_flag = 0 (for land and sun-glint observation)

If the data are to be compared with other XCO₂ data for which vertical profile information is available (e.g. inverse modeling, comparison to models, comparison to measured profiles), the column averaging kernels should be used. Here, it should be noted, that **the column averaging kernels are to be applied to layer sub-columns (m-2)**, as these are the quantities directly retrieved by the RemoTeC algorithm.

For model comparisons, the retrieved XCO₂ should be compared to $[VCO_2]_{\text{model}}/[VAIR]_{\text{model}}$, where $[VAIR]_{\text{model}}$ is the total dry air column provided by the model and $[VCO_2]_{\text{model}}$ is the model total CO₂ column after applying the column averaging kernel, viz.:

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 25
		Version 4.0
		15. August 2023

$$[VCO_2]_{\text{model}}' = [VCO_2]_{\text{prior}} + \mathbf{a}^T (\mathbf{x}_{\text{model}} - \mathbf{x}_{\text{prior}})$$

where $[VCO_2]_{\text{prior}}$ is the prior CO₂ total column used in the retrieval, $\mathbf{x}_{\text{model}}$ is the vertical CO₂ profile from the model (as sub-columns) and $\mathbf{x}_{\text{prior}}$ is the prior vertical profile from the retrieval. For application of the column averaging kernel, the model vertical profile should be re-calculated on the vertical grid of the retrieval (preferred) or the averaging kernel has to be interpolated to the vertical grid of the model.

5.4 Tools for Reading the Data

The data are stored in NetCDF format, which can be read with standard tools in the common programming languages (IDL, Matlab, Python, Fortran90, C++, etc).

 GHG-CCI+ project	ESA Climate Change Initiative “Plus” (CCI+) Product User Guide (PUG) XCO₂ GOSAT-2 SRON Full-Physics (CO₂_GO₂_SRFP) for the Essential Climate Variable (ECV) Greenhouse Gases (GHG)	Page 26
		Version 4.0
		15. August 2023

6. References

/Barr et al. 2023a/ A.G Barr and the GHG-CCI group at SRON: ESA Climate Change Initiative “Plus” (CCI+) ATBD (ATBDv4) for the RemoTeC CO₂ GOSAT-2 Data Product CO₂_GO₂_SRFP v2.0.2 for the Essential Climate Variable (ECV) Greenhouse Gases (GHG), ESA GHG CCI+, 2023.

/Barr et al. 2023b/ A.G Barr and the GHG-CCI+ group at SRON: ESA Climate Change Initiative “Plus” (CCI+) End-to-End ECV Uncertainty Budget Version 2.0.2 for the RemoTeC CO₂ GOSAT-2 Data Product CO₂_GO₂_SRFP for the Essential Climate Variable (ECV) Greenhouse Gases (GHG), ESA GHG CCI+, 2023.

/Laughner et al., 2021/ Laughner, J., Andrews, A., Roche, S., Kiel, M., and Toon, G. ginput v1.0.10: GGG2020 prior profile software (Version 1.0.10), <https://doi.org/10.22002/D1.1944>, 2021.

/Nakajima et al., 2017/ M. Nakajima, Hiroshi Suto, K. Yotsumoto, K. Shiomi, T. Hirabayashi, "Fourier transform spectrometer on GOSAT and GOSAT-2," Proc. SPIE 10563, International Conference on Space Optics — ICSO 2014, 105634O (17 November 2017); <https://doi.org/10.1117/12.2304062>.

/Toon et al., 2009/ Toon, G., Blavier, J.F., Washenfelder, R., Wunch, D., Keppel-Aleks, G., Wennberg, P., Connor, B., Sherlock, V., Griffith, D., Deutscher, N. and Notholt, J., 2009, April. Total column carbon observing network (TCCON). In Fourier Transform Spectroscopy (p. JMA3). Optical Society of America.

/Wunch et al., 2015/ Wunch, D., Toon, G.C., Sherlock, V., Deutscher, N.M., Liu, X., Feist, D.G., Wennberg, P.O., The Total Carbon Column Observing Network's GGG2014 Data Version. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA (available at: [doi:10.14291/tccon.ggg2014.documentation.R0/1221662](https://doi.org/10.14291/tccon.ggg2014.documentation.R0/1221662)), 2015.

*** end of document